

THE NOBEL PRIZE IN
CHEMISTRY 2008



Green Fluorescent Protein

Discovery of Green Fluorescent Protein (GFP)
(Nobel Lecture)**

Osamu Shimomura*

aequorin · bioluminescence · fluorescent proteins ·
Nobel Lecture

1. Biography

A Wartime Childhood

I was born on August 27, 1928 in the town of Fukuchiyama, Kyoto-Fu, Japan. My father Chikara was an army captain of the Fukuchiyama regiment. In the spring of 1933, my father took a post in Manchuria, which was under Japanese occupation. Because there were some insurgents in the Manchurian area, my mother Yukie, my younger brother Sadamu, and I chose to stay in Japan and moved to Sasebo, Nagasaki Prefecture, where we lived with my grandmother, Tsuki Shimomura. After the insurgents were wiped out, we moved to Renzankan, Manchuria in March 1935. A sister, Toshiko, was born here, but she died of pneumonia one year after her birth. I remember my mother's deep sorrow after Toshiko's death.

In early 1938, my father was transferred to an army position near the Soviet border. The rest of my family moved back to Sasebo. Mother soon departed to Manchuria again to live with my father, and Sadamu and I were in the custody of grandmother Tsuki for about one year. Because I was the elder son, Tsuki raised me carefully and strictly as the heir of the Shimomura family. Since I was physically not very strong, she tried to feed me various nutritious foods like omelet with ground beef and soybean milk. Grandmother was very strict about manners and etiquette. I always had to keep a good posture in her presence. She often said, "*the samurai betrays no weakness when starving.*" After I bathed, she would check behind my ears and neck for dirt. If she found any, she would say it would be ignominious to be dirty when I was beheaded (it is sometimes honorable for a samurai to do hara-kiri and then be beheaded). I knew she was talking about the importance of readiness, but it was a little scary. A year later, father took mother back home and stayed only briefly (Figure 1).

In April 1941, I entered the seventh grade at Sasebo middle school. Japanese militarism was maximal at the time. We had to perform military exercises once or twice a week under the direction of the officers attached to the school. Later that year, my father was transferred to Osaka, and after the Pacific War began on December 8, we joined my father there. At school in Osaka, my teacher happened to be Shizuo Ito, a well-known poet from Isahaya. I contracted tuberculosis and was often tired; although I studied hard, my class rank



Figure 1. Together with my father who returned briefly from Manchuria in 1939. Front row from left: my mother Yukie, grandmother Tsuki, grandfather Kosaburo, father Chikara, younger brother Sadamu, and myself behind Sadamu, together with my uncle Eijiro Sata's family in the back row.

was poor. I was interested in making model airplanes at the time. Once I built a plane based on an article in a scientific magazine, and I exhibited it at a large department store. My sister, Setsuko, was born, and I had to babysit occasionally.

When I was in the ninth grade, the war became clearly unfavorable to Japan, and my father was sent to Thailand. At school, missing lectures became common and military exercises increased. Once, on an overnight march to Mt. Ikoma, I saw many tiny irregular spots of light on the road. Now I think they were probably luminous earthworms crushed by shoes. In the summer of 1944, my father wrote us to move to the countryside to avoid American bombings. My mother chose to move us to her parents Fujiyama's house in Isayaha near Nagasaki. That September, on my first day of school (tenth grade), the teacher told us, "*This class is mobilized to the Omura Naval Aircraft Arsenal immediately. Other classes are mobilized to the Mitsubishi shipyard and arms factory in Nagasaki.*" Because the National General Mobilization Law

[*] Prof. O. Shimomura
Marine Biological Laboratory
Woods Hole, MA 02543 (USA)
E-mail: oshimomura@mbl.edu

[**] Copyright© The Nobel Foundation 2008. We thank the Nobel Foundation, Stockholm, for permission to print this lecture.

had been activated earlier, I expected to be mobilized sooner or later, but I didn't think it would be on the first day of school. From that day on, we did not listen to lectures or study at school: we worked.

Two months after I started working at the Omura Naval Aircraft Arsenal, the arsenal was destroyed by bombing. That day, when the air-raid siren sounded, some students took shelter in an underground bunker and the rest of us ran to the edge of the airfield, then waited in a ditch. Soon I saw a formation of more than 20 B-29 bombers coming from the west, and the bombing started. A stray bomb fell nearby and showered us with sand and gravel. The bombing destroyed most of the facilities, but I saw that one hangar was still standing and some people were trying to pull a fighter plane out of it. We rushed there to help them. However, an enemy plane was ahead in the smoke-covered sky. At the moment we reached the hangar, we were showered with numerous incendiary bombs. I heard someone shouting an order to take refuge. We ran between bombs burning with white flames; I saw a person who had been hit on the shoulder running with one arm dangling.

After the air raid, we learned that some of the students had died in the underground bunker, and that our dormitory had burned down. Within a month, several wooden factory buildings were built between the hills near Isahaya, and we were ordered to work there. I attended the factory every day, even before it was completed; if I found nothing to do I would often lie down in a sweet-potato field nearby and watch large formations of B-29 bombers going east, high above Mt. Tara-dake. It was beautiful to see the shining silver B-29s against the background of blue sky. Then, after about 10 minutes, I would see black smoke in the Ohmuta industrial area on the opposite shore of the Ariake Sea, and I could only imagine the scene of carnage over there.

The new factory was for the repair of fighter engines. My job was to smooth the face joining the crank case to the cylinder. My class graduated in March 1945, at the factory, without a graduation ceremony or diplomas. I was 16 years old. After graduation, the student mobilization continued.

The Nagasaki Atomic Bomb and the End of the War

On August 6, 1945, news reports informed us that the city of Hiroshima had been completely destroyed by a new type of bomb—we didn't know what kind. Three days later, shortly before 11 a.m., a siren sounded at the Isahaya factory, notifying us of an air raid. As usual, rather than going into a bunker, I went to the top of a nearby hill with a couple of friends and looked at the sky. We saw a single B-29 going from north to south towards Nagasaki, about 15 km away. I thought that its course was unusual. The B-29 dropped two or three parachutes and I heard sporadic gunshots. Watching carefully, I saw no people attached to the parachutes. Within a few minutes, another B-29 followed the first one, and a siren sounded the "all clear" signal. We returned to our factory building.

At the moment I sat down on my work stool, a powerful flash of light came through the small windows. We were

blinded for about 30 seconds. Then, about 40 seconds after the flash, a loud sound and sudden change of air pressure followed. We were sure there was a huge explosion somewhere, but we didn't know where. The sky was rapidly filling with dark clouds, and when I left the factory to walk home, about three miles away, a drizzling rain started. It was black rain. By the time I arrived home, my white shirt had turned gray. My grandmother quickly readied a bath for me. That bath might have saved me from the ill effects of the strong radiation that presumably existed in the black rain.

The next morning, a technical officer told us that the parachutes we had seen the day before contained measurement instruments and a transmitter. He also mentioned that there was serious damage in Nagasaki, but the details were unknown. The chief of the factory organized a rescue party. We tried to enter Nagasaki, but could not because the roads and the railroad were impassable. Later that afternoon, the railroad was opened to Michinoo, near Nagasaki station, and rescuers began to transport injured people to Isahaya and other cities.

On August 15, in a radio broadcast, Emperor Hirohito declared unconditional surrender. This was the first time that most Japanese citizens had heard the emperor's voice. I think there was a widespread feeling of relief, and also fear of an uncertain future.

Many years passed before we had detailed information about the atomic bombs that were dropped on Hiroshima and Nagasaki. The Nagasaki bomb was a different type and far more powerful than the Hiroshima bomb. Even if the use of the Hiroshima bomb was justifiable in order to precipitate an end to the war, the bomb dropped on Nagasaki three days later was clearly a test of new arms. It cannot be justified.

The student mobilization ended, but I was not sure if my school record would allow me to enter a college. To receive some guidance, I visited my Isahaya school a week after the end of the war. Attached to the main gate were several sheets of large paper, on which numerous names were written. About half of the names were crossed off. Apparently the school was being used to accommodate the injured people of Nagasaki, who were listed on the papers. In the exercise field, several men were slowly strolling under the strong summer sun amid the noisy shrill of cicada. All were half-naked. When I looked closer, I saw that their skin was covered with something black with white specks. I thought it was a black medicine (I later found out it was dried blood). The white specks were maggots that had hatched on the human flesh. But that was not the most shocking sight. At the front of the gate, bodies covered with straw mats were stacked on a cart, probably to be taken to a crematory. Two people with a stretcher, a body on it, were coming from the school building. At that moment I noticed two half-naked people standing near the fence. They had been watching the loading of the bodies, a process that might happen to them in a few days. I felt as if I were seeing ghosts. My brain froze, the shrill of cicada faded, and my senses vanished. I think the mental shock I had at that sight, when I was 16 years old, had a certain permanent effect on me. I have regretted for a long time that I did not speak to these people; probably I didn't have the courage at that time.

Nagasaki Pharmacy College

I tried to enter three different colleges in 1946 and 1947, but all rejected me. I didn't have a strong school record to assist me, because I hadn't studied even one day at the Isahaya school from which I graduated. I then heard that the Nagasaki Pharmacy College, a part of Nagasaki Medical College, was readying a temporary campus at a vacated military barracks near my home. (Both colleges had been completely destroyed by the atomic bomb.) I was admitted to the pharmacy college in April 1948. Although I was not planning to be a pharmacist, I didn't have any other choice under the circumstances. My grandmother presented me with a suit of silk clothes upon my entrance to college. There was no cloth to buy at the time, but she had a mulberry field; so she raised silkworms, reeled silk thread off the cocoons, dyed the thread, hand-wove the thread into cloth by herself, and then tailored the clothes.

Most of the pharmacy students lived in the dormitory, and they were always hungry due to an extreme shortage of food. I lived at home, and we owned farm land, so I was very lucky concerning food. Despite great effort by the professors, the resources and equipment at the pharmaceutical college were poor. We had a lot of difficulty carrying out experiments. Instruction in analytical chemistry and physical chemistry seemed fine, but I learned very little organic chemistry. In experiments on organic synthesis, we often set fire to solvents, because of poor technique and poor glassware, but many of us were experienced firefighters due to growing up in wartime, and we had no problem putting the fires out.

My interest in chemical experiments developed but, due to equipment limitations, the only experiments I could carry out were inorganic ionic reactions. With the permission of Professor Shungo Yasunaga (Figure 2, left) of the Pharma-




Figure 2. My three mentors. From left: Professor Shungo Yasunaga (1911–1959), Nagasaki University; Professor Yoshimasa Hirata (1915–2000), Nagoya University; and Professor Frank H. Johnson (1908–1990), Princeton University.

ceutical Analysis Laboratory, I prepared many glass capillaries 1 mm in diameter, and packed them with alumina powder. When a small amount of a mixed solution of metallic ions was applied to the tip of the capillary and then it was dipped into a developing reagent, the reagent quickly rose up by capillary action. In the process, metallic ions were separated, and colored bands appeared corresponding to the ions separated. It was a kind of chromatography. With Prof. Yasunaga's permission, I brought back a small amount of various chemicals to my home, and studied the conditions

of separation in detail. The results were reported in the *Journal of the Pharmaceutical Society of Japan* a few years later as my first paper.^[1]

In March 1951, I graduated from Nagasaki Pharmacy School at the top of my class. The school was reorganized as the Department of Pharmacy, Nagasaki University. Professor Yasunaga offered me a job there as an assistant in the analytical chemistry laboratory for students, and I took it. When I had spare time, I continued the chromatography experiments I had started when I was a student. I became interested in classical music after I heard an LP record somewhere. However, one LP cost nearly a month's salary. Somehow, probably with my parents' financial help, I managed to buy a record player and speakers, and I assembled an amplifier out of glass vacuum tubes, condensers, and resistors.

Professor Yasunaga was a gentle and very kind person, and trusted by people. After I had worked for him for four years, he obtained a leave of absence with pay for me to study for one year. Moreover, he offered to introduce me to Professor Fujio Egami of Nagoya University, a well-known molecular biologist. We took the train to Nagoya, but unfortunately we found Professor Egami wasn't at the university that day. Professor Yasunaga visited another person, Professor Yoshimasa Hirata (Figure 2, center), an organic chemist. We chatted a few minutes. As we were leaving, Professor Hirata said to me, "Come to my lab. You may start at any time." This was surprising, because we had just met. I didn't know much about molecular biology or organic chemistry, so it didn't matter to me which specialty I would study. I thought Professor Hirata's words might be the direction given by heaven, and I decided to go to his lab. It seems that this decision determined my future, directing me to the studies of bioluminescence, aequorin, and green fluorescent protein (GFP).

The Hirata Laboratory and *Cypridina* Luciferin

I enrolled in the Hirata laboratory, Department of Science, Nagoya University, as a research student in April 1955. The Hirata lab was a wonderful place, with a splendid atmosphere. Nobody taught me anything, but I learned much by watching other people and by independent study.

On my first day, Professor Hirata brought out a large vacuum desiccator and said, "This contains dried *Cypridina*." He explained to me that *Cypridina*, a small crustacean common in shallow coastal waters of Japan, emits light with an organic compound called luciferin and an enzyme luciferase. He told me that luciferin is extremely unstable and rapidly decomposes in the presence of oxygen, and also that Professor Newton Harvey of Princeton University had been trying to purify the luciferin for the past 20 years, but had been unsuccessful. Professor Hirata asked, "Could you purify and crystallize *Cypridina* luciferin for the purpose of structure determination?" At the time, crystallization was the only practical way to prove the purity of a substance. Professor Hirata also told me that he could not give this project to a student pursuing a degree, because the outcome was so

uncertain. I clearly understood the difficulty of the work. Since I was there for study, not for a degree, I replied, "I would like to do my best."

This did indeed turn out to be a very difficult crystallization. It took me ten months of extremely hard work to extract, purify, and crystallize the luciferin. When I finally succeeded, I was so happy I couldn't sleep for three days. Since the end of the war, my life had been dark, but this gave me hope for my future. Probably the greatest reward I gained was self-confidence; I learned that any difficult problem can be solved by great effort. With the crystallization accomplished, my stay in Nagoya was extended one year to study the structure of *Cypridina* luciferin (Figure 3). Our first paper on *Cypridina* luciferin was published in 1957, although the chromophore structure of luciferin remained to be elucidated.



Figure 3. The author with a close friend and collaborator Toshio Goto (right) in 1956, shortly after crystallizing *Cypridina* luciferin, behind a counter-current distribution apparatus.

To America

In the spring of 1959, I received a letter from Dr. Frank Johnson (Figure 2, right) of Princeton University inviting me to work at his laboratory. When Professor Hirata heard about my plan to go to Princeton, he awarded me a doctoral degree for my *Cypridina* work, even though I wasn't enrolled as a doctoral student. Prof. Hirata knew having a doctorate would double my salary at Princeton. I was completely surprised, and I accepted his offer with thanks. I applied for and received a Fulbright travel grant, which provided me with various experiences otherwise unavailable.

On August 4, 1960, I and Akemi Okubo got married in a traditional way, arranged by a match-maker. Akemi was a graduate of the Pharmacy Department where I worked. On August 27, I left Yokohama on a ship bound for Seattle, along with more than 200 other Fulbright fellows and students, but Akemi could not accompany me due to a visa problem. The day was my 32nd birthday. Since that voyage was the last Pacific cruise of the ship Hikawa-maru, the pier was filled with people (Figure 4). Thousands of colored tapes connected people on the boat with well-wishers on the pier. I will never forget the scene when the ship started to move, and the tapes broke and then fell. After traveling by rail across the United




Figure 4. The ship Hikawa-maru leaving Yokohama on August 27, 1960.

States, I arrived at Princeton on September 17, 1960, and stayed at Dr. Johnson's house that night. The next morning, Dr. Johnson offered to help me find an apartment. We saw a newspaper advertisement for a room for rent, and we stopped at the house. A man responded to the doorbell, but he quickly shut the door when he saw my face. It was a clear case of racial discrimination that I rarely encountered.

On my first visit to Dr. Johnson's office, he took out a small vial containing white powder, and said "This is the freeze-dried light organs of the luminous jellyfish *Aequorea*, and it should emit light when mixed with water." We went into a dark room and tested it, but we could not see any light. However, he enthusiastically explained to me that *Aequorea* were very abundant in Friday Harbor, Washington State, and that they were brilliantly luminous. Then he asked if I would like to study the bioluminescence of this jellyfish. I answered, "I will be glad to do it." Thus, we decided to go to Friday Harbor the following summer.

During my first few months in Princeton, I extracted and purified luciferase from dried *Cypridina*. I was surprised at the large quantity of dried *Cypridina* that was stocked in the Princeton biology department, which included sealed bottles dated 1928 and a sealed tin can bearing the name of Sakyo Kanda, a pioneer in the study of bioluminescence. There was also a large amount that had been collected for intended military use during the war, and seized by the US Navy at the end of war and then donated to Professor Harvey's lab. Dried *Cypridina*, even if it is many decades old, luminesces when crushed and mixed with water.

Miss Yo Saiga arrived in December 1960 to work as our research assistant, and my wife, Akemi, arrived the next month, just after the inauguration of President John F. Kennedy, after a delay of four months due to a visa problem.

On June 23, 1961, Dr. Johnson, my wife, Miss Saiga, and I departed for Friday Harbor. Dr. Johnson had purchased a new Plymouth station wagon for the trip. We loaded it with a large photometer, other instruments and chemicals needed for research, and suitcases for four people on the roof. It took us seven days to cross the continent, with Dr. Johnson driving the whole way. Upon arrival, we met our host, Dr. Robert Fernald, at the Friday Harbor Laboratories, University of Washington.

Friday Harbor is on San Juan Island, which lies east of the city of Victoria on Vancouver Island. The scenery of San Juan Island was splendid. The sea was clean and beautiful. At low tide, there were colorful sea urchins and starfish in various sizes scattered on rocks, and some abalones. There were also abundant fish. We could easily get rock fish of about 30 cm long using simple lures from a boat or rocky shore. We often ate them as sashimi. Indeed, Friday Harbor at the time was a kind of paradise for us.

For research space, Dr. Fernald assigned us part of a large laboratory. There were three other scientists in the room, and one of them was Dr. Dixy Lee Ray, a professor at the University of Washington at the time, and later chairperson of the US Atomic Energy Commission and governor of Washington State. She was always accompanied by her dog, even though dogs were prohibited in Friday Harbor Laboratory by a state law. She declared that the animal was her assistant.

Our research material, *Aequorea*, was really abundant. A constant stream of floating jellyfish passed along the side of the lab dock every morning and evening, riding with the tidal current. I describe our methods for collecting and extracting the luminescent material from *Aequorea* in my Nobel lecture. We brought the crude extract back to Princeton in September 1961, where we began the process of purifying it. In February 1962, we obtained about 5 mg of nearly pure luminescent substance. It was a protein, and we named it aequorin. Aequorin attracted wide attention as a unique protein that emits light in the presence of calcium ions, even in the absence of oxygen. Aequorin was the first photoprotein ever discovered. During the column chromatography of aequorin, we found a trace of protein that showed green fluorescence, which eluted sooner than aequorin. We also purified that protein, which is now called green fluorescent protein (GFP).

In August 1962, I visited Bermuda to study the famous fireworm *Odontosyllis enopla* at the Bermuda Biological Station. The fireworms are very small, but they show a spectacular bioluminescence display that is correlated with the lunar cycle. The luminescence display takes place during a period of several days following a full moon. It begins about one hour after sunset and lasts for only 10 minutes. It begins with the sudden appearance of a swarm of brilliantly luminescent females (about 2 cm long) at the surface of the water, each worm quickly moving in a tight circle. Within a few seconds, numerous brightly luminescent males (about 1 cm long) appear, and they dart toward the females from all directions, attracted to the light. I collected the fireworm, and studied the properties of its luciferin.

Due to my US visa expiring in 1963, I was expected to return to my position in the Pharmacy Department at Nagasaki University, from which I was on leave. However, I received through Professor Hirata an offer of the position of associate professor at the Water Science Institute, Nagoya University, with the understanding that I could continue my research on bioluminescence. I took the offer. In September 1963, I began working under Professor Tadashi Koyama at the Water Science Institute. A few months later, Professor Hirata asked me to assist his graduate student, Yoshito Kishi (later a professor at Harvard University) to determine the structure of *Cypridina* luciferin. In the summer of 1964, Kishi

and I went to Setoda in the Inland Sea to do mass collection of *Cypridina*, taking some ten students with us. From frozen *Cypridina*, we extracted luciferin and then purified and crystallized the luciferin. With that material, Kishi successfully determined the structure of the luciferin the next year.

In February 1965, I went to New Zealand to study two kinds of bioluminescent organisms: the cave worm *Arachnocampa* and the freshwater limpet *Latia*. Soon afterwards, I decided to go back to Dr. Johnson's lab at Princeton University, because I realized I did not have sufficient ability to do top-level work in two unrelated fields—earth science and bioluminescence. I wanted to study and clarify the chemical mechanism of aequorin bioluminescence, since some people doubted the existence of a photoprotein like aequorin. In December 1965, I returned to Princeton with my wife and our one-year-old son Tsutomu.

Since the early 1950s, it had been believed that calcium ions play important roles in living bodies, but there had been no way to demonstrate this experimentally. In 1967, however, Ellis Ridgway and Christopher Ashley at the University of Oregon proved experimentally the involvement of calcium ions in the contraction of muscles using aequorin as an indicator. As the usefulness of aequorin as a calcium probe increased in the fields of biology and physiology, I wanted to clarify the mechanism of aequorin luminescence to assist in its use.

I already knew that the aequorin luminescence was caused by an intramolecular reaction that takes place in the protein molecule, which would be difficult to study. But I decided to do my best. It turned out to be an unexpectedly large undertaking. It took about 12 years to obtain a structural model of aequorin, as detailed in my Nobel lecture. We needed a large amount of aequorin to carry this study out, so we returned to Friday Harbor every summer for more than 10 years, where we collected and processed about 3000 jellyfish a day. My staff, my wife, my children, and some local students that we hired were of great assistance in the collection and processing of *Aequorea*. Dr. Johnson devised a “jellyfish-cutting machine” to speed up the process of cutting the ring from the animals. In 1972, back at Princeton, we succeeded in determining the structure of AF350, a part of the aequorin chromophore. By 1978, we had achieved a general understanding of the aequorin luminescence reaction.

Between 1965 and 1978, in addition to my work with aequorin, I also did research on the bioluminescence of various luminous organisms including the limpet *Latia*, the krill *Meganctiphanes*, the worm *Chaetopterus*, the firefly squid *Watasenia*, various coelenterates, and luminous bacteria. I also studied the properties of GFP, and resolved a controversy regarding the role of a dioxetane intermediate in the luminescence reaction of firefly luciferin. Those matters are detailed in my book.^[2]

To the Marine Biological Laboratory, Woods Hole

Dr. Johnson retired from Princeton in 1977, and I decided to move to a marine laboratory. Before leaving Princeton, I elucidated the chromophore of GFP.^[3] I also performed

research on dinoflagellate luciferin and luminous scale worms. In 1981, with the kind arrangement made by Dr. Woodland Hastings of Harvard University and Dr. Benjamin Kaminer of Boston University, I moved to the Marine Biological Laboratory (MBL) in Woods Hole, Massachusetts, where I was a Senior Scientist. My wife, Akemi, worked at the MBL as my research assistant. I also accepted an adjunct professorship at Boston University Medical School.

Established in 1888, the MBL is the oldest marine laboratory in the western hemisphere. Over the years, many Japanese scientists have performed research at the MBL, including historic names such as Shosaburo Watase, Umeko Tsuda, Hideyo Noguchi, Sakyo Kanda, Katsuma Dan, Sachio Hiramoto, and Shinya Inoué. At the MBL, I continued my work to extract, purify, and study the luminescent substance of many organisms, including the millipede *Luminodesmus*, the brittle star *Ophiopsila*, and several types of luminous mushrooms.

Since 1975, aequorin had become widely used among cell biologists and physiologists as an excellent calcium probe, and its applications peaked around 1985. Since I was practically the only source of aequorin in this time period, I sent out several hundred aequorin samples in response to requests from investigators all over the world. The cDNA of aequorin was cloned and recombinant aequorin was made in 1985, but the patent owners of recombinant aequorin, the University of Georgia and Chisso Corporation of Japan, were not sure about expanding the use of aequorin, thus the general use of recombinant aequorin was delayed. In 1988, in collaboration with Professor Yoshito Kishi of Harvard, we succeeded in making various aequorins that had different calcium sensitivities and properties. Those aequorins contained various derivatives of coelenterazin instead of coelenterazine in their molecules, and were called semisynthetic aequorins. In 1995, I undertook work on the X-ray structure of aequorin, and the three-dimensional structure of aequorin was obtained in 2000.^[4]

In 1992, the cDNA of GFP was cloned by Dr. Douglas Prasher, who was then at Woods Hole Oceanographic Institution. At that time, however, it was commonly believed that expressing the cDNA in living organisms would not produce fluorescent GFP, because the formation of its chromophore requires condensation and dehydrogenation reactions, which are not expected to occur spontaneously. In 1994, however, Dr. Martin Chalfie of Columbia University tried to express the cDNA in *E. coli* and a nematode worm, and he and his colleagues unexpectedly observed the fluorescence of the expressed GFP. The results suggested GFP, and other proteins linked to GFP, could be expressed in living organisms to observe their behavior. Dr. Chalfie's work attracted the interest of many people, triggering rapid progress in the applications of GFP. Dr. Roger Tsien of the University of California, San Diego, engineered GFP by modifying the amino acid residues surrounding the chromophore, producing many different fluorescent proteins that emit various colors—from blue to red. Today, GFP is widely used as a fluorescent marker of protein molecules and cells, and it is an essential tool in the study of biology, physiology, and medicine. GFP has been used also in other fields, such as

the detection of cadmium, zinc, and explosive TNT and fumes.^[5] The range of applications of the fluorescent proteins is beyond imagination.

After My Retirement

I retired from the Marine Biological Laboratory in 2001. Because I wanted to do some more experiments and also I was still supplying aequorin samples to people upon request, I moved all my laboratory equipment and chemicals to my home, where I set up my photoprotein Laboratory. My retirement symposium, "GFP and Aequorin," was held at the MBL's Lillie Auditorium on July 27, 2002, through the kind arrangement of Dr. Shinya Inoué, a world authority on microscopy. Among the attendees were Martin Chalfie and Roger Tsien. I started to write a book for the next generation of students who want to explore the chemistry of bioluminescence. The book "*Bioluminescence: Chemical Principles and Methods*" was published by World Scientific Press in 2006. The book contains a comprehensive overview and chemical information on all known bioluminescence systems, comprising 35 different types of bioluminescent organisms.

The summer of 2004 was splendid. In July, I received the Pearse Prize from the Royal Microscopical Society for the discovery of GFP. In early August, I gave an invited lecture at the International Bioluminescence and Chemiluminescence Symposium. At the end of August, I attended the symposium "Calcium-Regulated Photoproteins and Green Fluorescent Protein", which was held in my honor at the Friday Harbor Laboratories as part of the lab's centennial celebration. At this symposium, almost all the well-known researchers in bioluminescence and related fields gathered from all over the world, and I was happy to see many of my old friends, although I was a little sad to see the sea at Friday Harbor was steadily being polluted. In 2006, I received the Asahi prize, one of the most prestigious prizes in Japan.

I would like to add a note on the jellyfish *Aequorea* at Friday Harbor. Between 1961 and 1988, we traveled to Friday Harbor and back to the East Coast 19 times (13 of which were road trips) and collected a total of about 850 000 *Aequorea* specimens to obtain aequorin for my research. The jellyfish were always abundant during that period. Mysteriously, however, they suddenly decreased in number after 1990, and it became very difficult to collect even a few. The cause for the drastic decrease could be pollution of the sea bed by crude oil spilled from the Exxon tanker Valdez in 1989, or it could be a natural cause. If the disappearance of the jellyfish had occurred 20 years earlier, we wouldn't know either the mechanism of the aequorin bioluminescence reaction or the chromophore of GFP.

2. The Nobel Lecture

Prologue

I discovered the green fluorescent protein GFP from the jellyfish *Aequorea aequorea* in 1961 as a by-product of the Ca-

sensitive photoprotein aequorin,^[6] and identified its chromophore in 1979.^[7] GFP was a beautiful protein, but it remained useless for the next 30 years after the discovery.

My story begins in 1945, the year the city of Nagasaki was destroyed by an atomic bomb and the Second World War ended. At that time I was a 16-year old high-school student, and I was working at a factory about 15 km northeast of Nagasaki. I watched the B-29 that carried the atomic bomb heading toward Nagasaki, then soon I was exposed to a blinding bright flash and a strong pressure wave that were caused by a gigantic explosion. I was lucky to survive the war. In the mess after the war, however, I could not find any school to attend. I idled for two years, then I learned that the pharmacy school of Nagasaki Medical College, which had been completely destroyed by the atomic bomb, was going to open a temporary campus near my home. I applied to the pharmacy school and was accepted. Although I didn't have any interest in pharmacy, it was the only way that I could have some education.

After graduating from the pharmacy school, I worked as a teaching assistant at the same school, which was reorganized as a part of Nagasaki University. My boss Professor Shungo Yasunaga was a gentle and very kind person. In 1955, when I had worked for four years on the job, he arranged for me a paid leave of absence for one year, and he sent me to Nagoya University, to study at the laboratory of Professor Yoshimasa Hirata.

Cypridina Luciferin

The research subject that Professor Hirata gave me was the bioluminescence of the crustacean ostracod *Cypridina hilgendorffii*. *Cypridina* emits blue light when its luciferin is oxidized in the presence of an enzyme luciferase and molecular oxygen (Figure 5). The luciferin had been studied for many years at Newton Harvey's laboratory at Princeton University,^[8] but it had never been completely purified, due to its extreme instability. Prof. Hirata wanted to determine the structure of the luciferin of *Cypridina*, and he asked me to purify the luciferin and to crystallize it, because crystallization




Figure 5. The ostracod *Cypridina hilgendorffii* freshly caught, and placed on a dark surface.

was the only way to confirm the purity of substances at the time.

Using 500 g of dried *Cypridina* (about 2.5 kg before drying), I began the extraction and purification of luciferin in an atmosphere of purified hydrogen using a large, specially made Soxhlet apparatus (Figure 6). After 5 days of day-and-




Figure 6. The apparatus used for extracting *Cypridina* luciferin.

night work, 500 g of dried *Cypridina* yielded about 2 mg of luciferin after purification. I tried to crystallize the purified luciferin, but all my efforts ended up with amorphous precipitates, and any leftover luciferin became useless through oxidation by the next morning. So I had to repeat the extraction and purification again and again. I worked very hard, and tried every method of crystallization that I could think of, without success. Ten months later, however, I finally found that the luciferin could be crystallized in a highly unusual solvent.^[9] The solvent I found was a high concentration of hydrochloric acid. Using the crystallized luciferin, we were able to determine the chemical structures of the luciferin and its oxidation products (Figure 7).^[10] Those data became essential later in the study of aequorin.

In 1959, Professor Frank Johnson invited me to work at his Princeton laboratory owing to my success in the *Cypridina*




Figure 7. Crystals of *Cypridina* luciferin (left), and the chemical structures of *Cypridina* luciferin and its luminescence reaction product, oxyluciferin (right).

work. In September 1960, shortly after my arrival at Princeton, Dr. Johnson asked me if I would be interested in studying the bioluminescence of the jellyfish *Aequorea*. I was strongly impressed by his description of the brilliant luminescence and the abundance of the jellyfish at Friday Harbor in the state of Washington. I agreed to study the jellyfish.

The Jellyfish *Aequorea* and the Photoprotein Aequorin

Early in the summer of 1961, we traveled from Princeton to Friday Harbor, driving 5000 kilometers. Friday Harbor was a quiet, peaceful small village at the time (Figure 8). The jellyfish were abundant in the water (Figure 9). At the University of Washington laboratory there, we carefully scooped up the jellyfish one by one using a shallow dip net. The light organs of *Aequorea aequorea* (Figure 10) are located along the edge of the umbrella (which we called a ring). The ring could be cut off with a pair of scissors, eliminating most of the unnecessary body part.

At the time, it was a common belief that the light of all bioluminescent organisms was produced by the reaction of



Figure 8. Friday Harbor, 1961. The University of Washington Laboratory is located on the opposite shore of the bay, to the left of the center in the picture. By 1980, about one third of the bay in the foreground had been turned into a marina.




Figure 9. The jellyfish *Aequorea aequorea* in nature.




Figure 10. Top view of a specimen of *Aequorea aequorea* in daylight at sea (left), and when stimulated in a darkroom (right).

luciferin and luciferase. Therefore, we tried to extract luciferin and luciferase from the rings of the jellyfish. We tried every method we could think of, but all our efforts failed. After only a few days of work, we ran out of ideas.

I was convinced that the cause of our failure was the luciferin–luciferase hypothesis that dominated our minds. I suggested to Dr. Johnson that we forget the idea of extracting luciferin and luciferase and, instead, try to extract a luminescent substance, whatever it might be. However, I was unable to convince him. Because of the disagreement on the experimental method, I started to work alone at one side of a table, while, on the other side, Dr. Johnson and his assistant continued their efforts to extract a luciferin. It was an awkward, uncomfortable situation.

Since the emission of light means the consumption (loss) of active bioluminescent substance, the extraction of bioluminescent substances from light organs must be performed under conditions that reversibly inhibits the luminescence reaction. Therefore, I tried to reversibly inhibit luminescence with various kinds of inhibitors of enzymes and proteins. I tried very hard, but nothing worked. I spent the next several days soul-searching, trying to find out something missing in my experiments and in my thought. I thought day and night. I often took a rowboat out to the middle of the bay to avoid interference from people. One afternoon, an idea suddenly struck me on the boat. It was a very simple idea: “The luminescence reaction probably involves a protein. If so, luminescence might be reversibly inhibited at a certain pH.”

I immediately went back to the lab and tested the luminescence of light organs at various pH values. I clearly saw luminescence at pH 7, 6, and 5, but not at pH 4. I ground the light organs in a pH 4 buffer, then filtered the mixture. The cell-free filtrate was nearly dark, but it regained luminescence when it was neutralized with sodium bicarbonate. The experiment showed that I could extract the luminescence substance, at least in principle (Figure 11).

But a big surprise came the next moment. When I threw the extract into a sink, the inside of the sink lit up with a bright blue flash. The overflow of an aquarium was flowing into the sink, so I figured out that seawater had caused the luminescence. Because the composition of seawater is known, I easily found out that Ca^{2+} ions activated the luminescence. The discovery of Ca^{2+} as the activator suggested that the luminescence material could be extracted utilizing the Ca-chelator EDTA (ethylenediaminetetraacetate), and we




Figure 11. The process that revealed Ca²⁺ to be the activator of luminescence reaction.



Figure 12. Procedure for extraction of aequorin and GFP.

devised an extraction method of the luminescent substance (Figure 12).

In the rest of the summer of 1961, we extracted the luminescent substance from about 10 000 jellyfish. After returning to Princeton, we purified the luminescent substance, and obtained a few milligrams of purified protein. The protein emitted blue light in the presence of a trace of Ca²⁺.

We named the protein aequorin.^[6] Aequorin was the first example of a photoprotein discovered.^[11] During the purification of aequorin, we found another protein that exhibited a bright green fluorescence. It was only in a trace amount, but we purified this protein too, and called it “green protein”. The protein was renamed “green fluorescent protein” by Morin and Hastings.^[12]

We wanted to understand the mechanism of the aequorin bioluminescence reaction, because it became clear in 1967 that aequorin was highly useful and important as a calcium probe in biological studies.^[13] First, we tried to isolate the light-emitting chromophore of aequorin. However, there was no way to extract the native chromophore,^[14,15] because any attempt to extract the chromophore always resulted in an intramolecular reaction of aequorin that triggered the emission of light, destroying the original chromophore. Indeed, the secret of light emission of *Aequorea* was well protected.

We nevertheless found that a fluorescent compound was formed when aequorin was denatured with urea in the presence of 2-mercaptoe-

thanol.^[14] We named this fluorescent compound AF-350, based on its absorption maximum at 350 nm. We decided to determine the structure of AF-350. However, to obtain the 1 mg of AF-350 needed for a single structural study of this compound, about 150 mg of purified aequorin was needed, and that meant we had to collect and extract at least 50 000 jellyfish. Considering that we would probably need several milligrams of AF-350, the structure determination was a huge undertaking for us.

In processing a large number of jellyfish to obtain a sufficient amount of AF-350, we found that cutting rings with a pair of scissors was too slow. To speed up the process, Dr. Johnson constructed a jellyfish cutting machine (Figure 13), which enabled one person to cut more than 600 rings per hour—10 times more than by hand.

We started to collect jellyfish at 6 a.m., and a part of our group began to cut off the rings at 8 a.m. We spent all afternoon extracting aequorin from the rings. Then, we collected more jellyfish in the evening (7 to 9 p.m.) for the next day (Figure 14). Our laboratory looked like a jellyfish factory (Figure 15), and was filled with the jellyfish smell.

After five years of hard work, we determined the chemical structure of AF-350 in 1972.^[16] The result was surprising. The structure of AF-350 contained the skeleton of a 2-amino-pyrazine (Figure 16) that was previously found in the oxidation products of *Cypridina* luciferin, although the side chains are different. This finding suggested a close relationship between the luminescence systems of *Aequorea* and *Cypridina*. Based on that information, we were able to determine the structure of the chromophore of aequorin to be coelenterazine (Figure 16). Eventually we elucidated that the luminescence reaction of aequorin takes place as shown in Figure 17.




Figure 13. The jellyfish ring cutting machine constructed by Frank H. Johnson in 1969. A specimen is placed on the black plexiglas platform and rotated to spread the edge of the umbrella. While rotating, the specimen is pushed toward the rotating blade (10-inch meat cutting blade) to cut off a 2–3 mm wide strip containing the light organs. The strip drops into a container below.



Figure 14. Left: Jellyfish collectors of the summer of 1974. From left, my wife Akemi, Dr. Chang, myself, Mrs. Chang, Mrs. Johnson, Dr. Johnson, Debby (a helper), my children, Tsutomu and Sachi. We collected 30–40 bucketfuls of jellyfish each day. Right: My family collecting jellyfish.



Figure 15. Scenes of jellyfish cutting by two assistants (left), and the extraction of aequorin (right).




Figure 16. The chemical structures of AF-350 (coelenteramine), coelenteramide (a product of the luminescence reaction of aequorin), and coelenterazine, compared with those of *Cypridina oxyluciferin* and luciferin.

Green Fluorescent Protein

In a live specimen of *Aequorea*, the light organs contain GFP in addition to aequorin, and the energy of the blue light




Figure 17. The luminescence and regeneration of aequorin. The photo-protein aequorin binds with two Ca^{2+} ions,^[17,18] and decomposes into coelenteramide, CO_2 , and apoaequorin accompanied by the emission of light (emission maximum at 465 nm). Apoaequorin can be regenerated into the original aequorin by incubation with coelenterazine in the presence of oxygen.

produced by the aequorin molecule is transferred to the GFP molecule, and GFP emits green light.^[19]

Although GFP is highly visible and readily crystallizable (Figure 18, left), the yield of GFP from the jellyfish was extremely low, much lower than that of aequorin. Therefore, to study GFP, we had to accumulate GFP little by little for many years while we studied the chemistry of aequorin luminescence. The amount of GFP we accumulated reached a sufficient amount to study this protein in 1979. Thus, we tried to find out the nature of the GFP chromophore by a series of experiments, using 100 mg of the protein in one experiment (Figure 18, right).

We first cut the molecule of GFP into small pieces of peptide by enzymic digestion. We isolated and purified the peptide that contained the chromophore, and then analyzed the structure of the chromophore. I was surprised when I




Figure 18. Left: Crystals of GFP (photo by Dr. Shinya Inoué); right: the process of isolating the chromophore of GFP.

measured the absorption spectrum of the peptide. The spectrum was nearly identical to that of a compound that I had synthesized in my study of the *Cypridina* luciferin 20 years earlier. Based on the spectral resemblance and some other properties, I could quickly identify the chromophore structure of GFP.^[7]

What I found is illustrated in Figure 19. Fluorescent proteins are usually a complex of a protein and a fluorescent compound (as shown in Figure 19 upper left). However, GFP was a very special fluorescent protein that contained a fluorescent chromophore within the protein molecule (as shown in Figure 19 upper right). The lower part of the figure




Figure 19. The structure of the chromophore in the molecules of GFP.

shows further details of GFP. GFP is a single peptide chain containing more than 200 amino acid residues. The chromophore was formed from 3 amino acid residues in the peptide chain by dehydration and dehydrogenation reactions. This finding was extremely important because it showed that the chromophore is a part of the peptide chain, and thus it opened up the possibility of cloning GFP. The chromophore structure was later confirmed by Cody et al. (1993).^[20]

When I found the chromophore of GFP in 1979, I thought I had done all I could do with GFP, and decided to terminate my work on GFP in order to concentrate my efforts in the study of bioluminescence, my lifework. Then a mysterious thing happened. The population of *Aequorea* in the Friday Harbor area drastically decreased after 1990, thus making it

practically impossible to prepare any new samples of natural aequorin or GFP. Fortunately, however, aequorin had been cloned by Inouye et al.^[21,22] and Prasher et al.^[23] and GFP was cloned by Prasher et al.^[24] thus making the natural proteins unessential. In 1994, GFP was successfully expressed in living organisms by Chalfie et al.^[25] and it was further developed into the present prosperous state by Roger Tsien.

Nowadays GFP and its homologues are indispensable in biomedical research, due to the fact that these proteins self-contain a fluorescent chromophore in their peptide chains and they can be expressed in living bodies. The identification of the fluorescent chromophore, however, depended on the GFP that had been accumulated for many years in our study of aequorin. Without the study of aequorin, the chromophore of GFP would have remained unknown and the flourishing of fluorescent proteins would not have occurred.

Received: April 24, 2009

Published online: July 3, 2009

- [1] "Chromatography of inorganic ions. I. Capillary microanalysis with alumina for cations": S. Yasunaga, O. Shimomura, *Yakugaku Zasshi* **1953**, 73, 1346–1350, in Japanese.
- [2] O. Shimomura, *Bioluminescence: Chemical Principles and Methods*, World Scientific Publishing, Hackensack, N.J., **2006**, p. 498.
- [3] "Structure of the chromophore of *Aequorea* green fluorescent protein": O. Shimomura, *FEBS Lett.* **1979**, 104, 220–222.
- [4] "The crystal structure of the photoprotein aequorin at 2.3 Å resolution": J. F. Head, S. Inouye, K. Teranishi, O. Shimomura, *Nature* **2000**, 405, 372–376.
- [5] M. Zimmer, *Glowing genes*. Prometheus, Amherst, **2005**, p. 221.
- [6] "Extraction, purification and properties of aequorin, a bioluminescent protein from the luminous hydromedusan, *Aequorea*": O. Shimomura, F. H. Johnson, Y. Saiga, *J. Cell. Comp. Physiol.* **1962**, 59, 223–239.
- [7] "Structure of the chromophore of *Aequorea* green fluorescent protein": O. Shimomura, *FEBS Lett.* **1979**, 104, 220–222.
- [8] E. N. Harvey, *Bioluminescence*, Academic Press, New York, **1952**.
- [9] "Crystalline *Cypridina* luciferin": O. Shimomura, T. Goto, Y. Hirata, *Bull. Chem. Soc. Jpn.* **1957**, 30, 929–933.
- [10] "Cypridina bioluminescence I: structure of *Cypridina* luciferin": Y. Kishi, T. Goto, Y. Hirata, O. Shimomura, F. H. Johnson, *Tetrahedron Lett.* **1966**, 7, 3427–3436.
- [11] "Bioluminescence in the sea: photoprotein systems": O. Shimomura, *Symp. Soc. Exp. Biol.* **1985**, 39, 351–372.
- [12] "Energy transfer in a bioluminescent system": J. G. Morin, J. W. Hastings, *J. Cell. Physiol.* **1971**, 77, 313–318.
- [13] "Calcium transients in single muscle fibers": E. B. Ridgway, C. C. Ashley, *Biochem. Biophys. Res. Commun.* **1967**, 29, 229–234.
- [14] "Properties of the bioluminescent protein aequorin": O. Shimomura, F. H. Johnson, *Biochemistry* **1969**, 8, 3991–3997.
- [15] "Mechanism of the luminescent intramolecular reaction of aequorin": O. Shimomura, F. H. Johnson, H. Morise, *Biochemistry* **1974**, 13, 3278–3286.
- [16] "Structure of the light-emitting moiety of aequorin": O. Shimomura, F. H. Johnson, *Biochemistry* **1972**, 11, 1602–1608.
- [17] "Luminescence of aequorin is triggered by the binding of two calcium ions": O. Shimomura, *Biochem. Biophys. Res. Commun.* **1995**, 211, 359–363.

- [18] "Titration of recombinant aequorin with calcium chloride": O. Shimomura, S. Inouye, *Biochem. Biophys. Res. Commun.* **1996**, *221*, 77–81.
- [19] "Intermolecular energy transfer in the bioluminescent system of *Aequorea*": H. Morise, O. Shimomura, F. H. Johnson, J. Winant, *Biochemistry* **1974**, *13*, 2656–2662.
- [20] "Chemical structure of the hexapeptide chromophore of the *Aequorea* green-fluorescent protein": C. W. Cody, D. C. Prasher, W. M. Westler, F. G. Prendergast, W. W. Ward, *Biochemistry* **1993**, *32*, 1212–1218.
- [21] "Cloning and sequence analysis of cDNA for the luminescent protein aequorin": S. Inouye, M. Noguchi, Y. Sakaki, Y. Takagi, T. Miyata, S. Iwanaga, T. Miyata, F. I. Tsuji, *Proc. Natl. Acad. Sci. USA* **1985**, *82*, 3154–3158.
- [22] "Expression of apoaequorin complementary DNA in *Escherichia coli*": S. Inouye, Y. Sakaki, T. Goto, F. I. Tsuji, *Biochemistry* **1986**, *25*, 8425–8429.
- [23] "Cloning and expression of the cDNA coding for aequorin, a bioluminescent calcium-binding protein": D. Prasher, R. O. McCann, M. J. Cormier, *Biochem. Biophys. Res. Commun.* **1985**, *126*, 1259–1268.
- [24] "Primary structure of the *Aequorea victoria* green fluorescent protein": D. C. Prasher, V. K. Eckenrode, W. W. Ward, F. G. Prendergast, M. J. Cormier, *Gene* **1992**, *111*, 229–233.
- [25] "Green fluorescent protein as a marker for gene expression": M. Chalfie, Y. Tu, G. Euskirchen, W. W. Ward, D. C. Prasher, *Science* **1994**, *263*, 802–805.
-